

GAPDH Loading Control Monoclonal Antibody (GA1R)

Product Details

Size	100 µg
Species Reactivity	Bacteria, Dog, Chicken, Hamster, Human, Insect, Mouse, Rabbit, Rat, Yeast
Published Species	Dog, Yeast, Rabbit, Rat, Pig, Bacteria, Human, Mouse, Chicken
Host/Isotope	Mouse / IgG1
Class	Monoclonal
Type	Antibody
Clone	GA1R
Conjugate	Unconjugated
Immunogen	Recombinant GAPDH
Form	Liquid
Concentration	1 mg/mL
Purification	Protein A
Storage buffer	PBS, pH 7.2
Contains	no preservative
Storage Conditions	Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.
RRID	AB_10977387

Applications	Tested Dilution	Publications
ELISA (ELISA)	Assay Dependent	-
Flow Cytometry (Flow)	2 µg/test	-
Immunocytochemistry (ICC)	1:20-1:200	60 Publications
Immunofluorescence (IF)	1:20-1:200	-
Immunohistochemistry (Paraffin) (IHC (P))	1:50-1:500	-
Western Blot (WB)	1:1000-1:10,000	152 Publications
Miscellaneous PubMed (Misc)	-	12 Publications
in situ PLA (PLA)	-	2 Publications


Product Specific Information

This antibody detects GAPDH from BL-21 bacteria, Sf9 insect, *Saccharomyces cerevisiae* (yeast), human, mouse, rat, rabbit, hamster, and chicken samples. MA5-15738 has been successfully used in Western blot, ICC, IF, IHC (P), FACS and ELISA. MA5-15738 may be stored at 4°C short term. Add 0.05% sodium azide if desired. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

Product Images For GAPDH Loading Control Monoclonal Antibody (GA1R)


GAPDH Loading Control Antibody (MA5-15738) in IF

Immunofluorescent analysis of GAPDH (green) in HeLa. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 5% normal goat serum (Product # 31873) for 15 minutes at room temperature. Cells were probed with a GAPDH monoclonal antibody (Product # MA5-15738) at a dilution of 1:100 for at least 1 hour at room temperature, washed with PBS, and incubated with DyLight 488 goat anti-mouse IgG secondary antibody (Product # 35502) at a dilution of 1:400 for 30 minutes at room temperature. Nuclei (blue) were stained with Hoechst 33342 dye (Product # 62249). Images were taken on a Thermo Scientific ArrayScan or ToxInsight Instrument at 20X magnification.




GAPDH Loading Control Antibody (MA5-15738) in IHC (P)

Immunohistochemistry analysis of GAPDH showing staining in the nucleus and cytoplasm of paraffin-embedded human breast carcinoma (right) compared with a negative control without primary antibody (left). To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 min. Following antigen retrieval, tissues were blocked in 3% H₂O₂-methanol for 15 min at room temperature, washed with ddH₂O and PBS, and then probed with a GAPDH monoclonal antibody (Product # MA5-15738) diluted in 3% BSA-PBS at a dilution of 1:200 overnight at 4°C in a humidified chamber. Tissues were washed extensively in PBST and detection was performed using an HRP-conjugated secondary antibody followed by colorimetric detection using a DAB kit. Tissues were counterstained with hematoxylin and dehydrated with ethanol and xylene to prep for mounting.



GAPDH Loading Control Antibody (MA5-15738) in WB

1:2000 (0.5 µg/mL) Ab dilution used in WB of 5 µg/lane tissue lysates from human (1), mouse (2), rat (3), rabbit (4), chicken (5), and hamster (6).



[View more figures on thermofisher.com](https://www.thermofisher.com)

Immunocytochemistry (60)

Frontiers in cellular neuroscience

Herpes Simplex Virus Type 1 Enhances Expression of the Synaptic Protein Arc for Its Own Benefit.

"MA5-15738 was used in Western Blotting to study the modulation of synaptic Arc expression by Herpes Simplex Virus Type 1 infection."

Authors: Acuña-Hinrichsen F, Muñoz M, Hott M, Martin C, Mancilla E, Salazar P, Leyton L, Zambrano A, Concha MI, Burgos PV, Otth C

Species
Human

Dilution
Not Cited

Year
2020

Cell death discovery

PEGylated liposome-encapsulated rhenium-188 radiopharmaceutical inhibits proliferation and epithelial-mesenchymal transition of human head and neck cancer cells in vivo with repeated therapy.

"MA5-15738 was used in Western Blotting to investigate the biological and pharmaceutical responses of high rhenium-188-liposomal dosage in vivo."

Authors: Chang CY, Chen CC, Lin LT, Chang CH, Chen LC, Wang HE, Lee TW, Lee YJ

Species
Mouse

Dilution
Not Cited

Year
2020

[View more ICC references on thermofisher.com](#)

Western Blot (152)

Frontiers in molecular neuroscience

The Enzymatic Core of the Parkinson's Disease-Associated Protein LRRK2 Impairs Mitochondrial Biogenesis in Aging Yeast.

"MA5-15738 was used in Western Blotting to study the effects of high levels of the enzymatic core of human leucine-rich repeat kinase 2 on mitochondrial biogenesis in ageing yeast."

Authors: Aufschnaiter A, Kohler V, Walter C, Tosal-Castano S, Habernig L, Wolinski H, Keller W, Vögtle FN, Büttner S

Species
Yeast

Dilution
Not Cited

Year
2020

Cell death discovery

FoxO1 interacts with transcription factor EB and differentially regulates mitochondrial uncoupling proteins via autophagy in adipocytes.

"MA5-15738 was used in western blot to elucidate how FoxO1 regulates mitochondrial uncoupling proteins"

Authors: Liu L, Tao Z, Zheng LD, Brooke JP, Smith CM, Liu D, Long YC, Cheng Z

Species
Mouse

Dilution
Not Cited

Year
2020

[View more WB references on thermofisher.com](#)

More applications with references on thermofisher.com

PLA (2) **Misc (12)**

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.