

F4/80 Monoclonal Antibody (BM8), APC, eBioscience™

Product Details	
Size	100 µg
Species Reactivity	Mouse
Published Species	Hamster, Fish, Mouse, Human
Host/Isotope	Rat / IgG2a, kappa
Recommended Isotype Control	Rat IgG2a kappa Isotype Control (eBR2a), APC, eBioscience™
Class	Monoclonal
Type	Antibody
Clone	BM8
Conjugate	APC
Form	Liquid
Concentration	0.2 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.1% gelatin
Contains	0.09% sodium azide
Storage Conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2784648

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	2 µg/test	289 Publications
Immunofluorescence (IF)	-	8 Publications
Immunohistochemistry (Frozen) (IHC (F))	-	2 Publications
Immunohistochemistry (IHC)	-	2 Publications
Immunohistochemistry (PFA fixed) (IHC (PFA))	-	1 Publication
Immunohistochemistry (Paraffin) (IHC (P))	-	1 Publication
Miscellaneous PubMed (Misc)	-	2 Publications

Product Specific Information

Description: The BM8 monoclonal antibody reacts with mouse F4/80 antigen, an approximately 160 kDa surface receptor. It belongs to the EGF-TM7 family of proteins. As such it contains seven EGF-like domains on its extracellular N-terminus, seven transmembrane spanning sequences, and an intracellular C-terminal domain showing homology to other TM7 superfamily members. The F4/80 antigen is expressed by a majority of mature macrophages and is one of the best markers for this population of cells. However, other cell types, such as peritoneal eosinophils, Langerhans cells, and some other dendritic cell subtypes, have been reported to express this antigen as well. Expression of F4/80 commences during early myeloid development in vivo and can be upregulated on BM cells stimulated in vitro with M-CSF. Some populations of macrophages, especially in the lymphoid microenvironment, may be devoid of F4/80.


Applications Reported: BM8 has been reported for use in flow cytometric analysis.

Applications Tested: The BM8 antibody has been tested by flow cytometric analysis of mouse resident peritoneal exudate cells. This can be used at less than or equal to 2 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Excitation: 633-647 nm; Emission: 660 nm; Laser: Red Laser.

Filtration: 0.2 µm post-manufacturing filtered.

Product Images For F4/80 Monoclonal Antibody (BM8), APC, eBioscience™



F4/80 Antibody (17-4801-82) in Flow

Staining of C57Bl/6 resident peritoneal exudate cells with Anti-Mouse CD11b PE (Product # 12-0112-82) and 1 µg of Rat IgG2a K Isotype Control APC (Product # 17-4321-81) (left) or 1 µg of Anti-Mouse F4/80 Antigen APC (right). Cells in the large scatter population were used for analysis.

[View more figures on thermofisher.com](http://thermofisher.com)

Flow Cytometry (289)

Frontiers in immunology

Prolonged Deleterious Influences of Chemotherapeutic Agent CPT-11 on Resident Peritoneal Macrophages and B1 Cells.

"Published figure using F4/80 monoclonal antibody (Product # 17-4801-82) in Flow Cytometry"

Authors: Bai WJ,Li CG,Zhang CC,Xu LH,Zeng QZ,Hu B,Hong Z,He XH,Ouyang DY

Species

Not Applicable

Dilution

Not Cited

Year

2019

Frontiers in immunology

p40 -Deficient Mice Exhibit Impaired Bacterial Clearance and Enhanced Pro-inflammatory Responses during *Salmonella enterica* serovar Typhimurium Infection.

"Published figure using F4/80 monoclonal antibody (Product # 17-4801-82) in Flow Cytometry"

Authors: Li Y,Lv M,Su C,Long S,Zhang W,Conway KL,Li W,Xavier RJ,Shi HN

Species

Not Applicable

Dilution

Not Cited

Year

2019

[View more Flow references on thermofisher.com](#)

Miscellaneous PubMed (2)

Oncotarget

Tumor-associated macrophages promote neuroblastoma via STAT3 phosphorylation and up-regulation of c-MYC.

"17-4801 was used in Flow cytometry/Cell sorting to indicate a novel role of tumour-associated macrophages as inducers of MYC expression in neuroblastomas lacking independent oncogene activation."

Authors: Hadjidaniel MD,Muthugounder S,Hung LT,Sheard MA,Shirinbak S,Chan RY,Nakata R,Borriello L,Malvar J,Kennedy RJ,Iwakura H,Akamizu T,Spoto R,Shimada H,DeClerck YA,Asgharzadeh S

Species

Mouse

Dilution

Not Cited

Year

2017

Cell metabolism

A macrophage NBR1-MEKK3 complex triggers JNK-mediated adipose tissue inflammation in obesity.

"Published figure using F4/80 monoclonal antibody (Product # 17-4801-82) in Miscellaneous"

Authors: Hernandez ED,Lee SJ,Kim JY,Duran A,Linares JF,Yajima T,Müller TD,Tschöp MH,Smith SR,Diaz-Meco MT,Moscat J

Species

Mouse
Not Applicable

Dilution

Not Cited
Not Cited

Year

2014

More applications with references on thermofisher.com

IF (8) IHC (2) IHC (PFA) (1) IHC (F) (2) IHC (P) (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.