

FITC anti-mouse CD95 (Fas) Antibody

Catalog# / Size	152605 / 25 µg 152606 / 100 µg
Clone	SA367H8
Other Names	Fas, ALPS1A, APO-1, APT1
Isotype	Mouse IgG1, κ
Description	CD95, also known as Fas, is an approximately 45 kD type I transmembrane protein belonging to the TNFR superfamily (TNFRSF6). The expression of CD95 has been described in various organs, such as thymus, spleen, liver, heart, lung and ovary. Upon ligand (FasL) binding, CD95 forms the death-inducing signaling complex (DISC) intracellularly and induce apoptosis of the cell. CD95-induced apoptosis plays important roles in development, as well as in maintaining peripheral tolerance of the immune system.

Product Details

Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Mouse Fas Transfectants
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.125 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm)
RRID	AB_2632900 (BioLegend Cat. No. 152605) AB_2632901 (BioLegend Cat. No. 152606)

Antigen Details

Structure	CD95 (Fas) is an approximately 45 kD type I transmembrane protein; cysteine-rich domain, death domain.
Distribution	Thymus, spleen, liver, heart, lung, and ovary.
Function	Induction of apoptosis.
Interaction	Caspase 8, caspase 10, CFLAR, FADD, PDCD6, SUMO1, and TNFRSF6.
Ligand/Receptor	FasL.
Biology Area	Apoptosis/Tumor Suppressors/Cell Death, Cell Biology, Immunology
Molecular Family	CD Molecules
Antigen References	1. Ogasawara J, et al. 1995. J. Exp. Med. 181:485. 2. Ogasawara J, et al. 1993. Nature. 364:806. 3. Nishimura Y, et al. 1995. J. Immunol. 154:4395.


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-mouse CD95 (Fas), PerCP/Cyanine5.5 anti-mouse CD95 (Fas), APC anti-mouse CD95 (Fas), Purified anti-mouse CD95 (Fas), Brilliant Violet 605™ anti-mouse CD95 (Fas), TotalSeq™-A0917 anti-mouse CD95 (Fas), TotalSeq™-C0917 anti-mouse CD95 (Fas)

Product Data



C57BL/6 thymocytes were stained with CD95 (clone SA367H8) FITC (filled histogram) or mouse IgG1, κ FITC isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587