

Datasheet: MCA822

Description:	MOUSE ANTI HUMAN CD126/IL-6R
Specificity:	CD126
Other names:	INTERLEUKIN-6 RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	B-R6
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been repo	This product has been reported to work in the following applications. This information is derived					
	from testing within our labo	from testing within our laboratories, peer-reviewed publications or personal communications from					
	the originators. Please refer to references indicated for further information. For general protocol						
	recommendations, please visit www.bio-rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			1/50		
	Immunohistology - Frozen	-					
	Immunohistology - Paraffin						
	ELISA	-					
	Immunoprecipitation	-					
	Western Blotting						
	Immunofluorescence	-					
	Where this antibody has no	t been tes	sted for use	in a particular technique	this does not necessarily		
	exclude its use in such proc	cedures. S	Suggested v	vorking dilutions are give	n as a guide only. It is		
	recommended that the user titrates the antibody for use in their own system using a				em using appropriate		
	negative/positive controls.		-				
	<u> </u>						
Target Species	Human						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by af	finity chro	matography	on Protein A from tissue	e culture supernatant		
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide						
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml						
Immunogen	CHO cells transfected with	cDNA of t	he IL-6 rece	eptor.			
External Database	UniProt:						

Ritoz Gono: 3570 ILGR Belated reagents RKD AB_2127915 Fusion Partners Spleen cellis from immunised BALB/c (Iffa Credo) mice were fused with the X63/Ag.8653 mouse myeloma cell line. Specificity Mouse anti Human CD126 antibody, clone B-R6 recognizes human interleukin-6 receptor subunit alpha, also known as CD126, IL-R6 or Membrane glycoprotein 80. Human CD126 is a 468 animo acid singipe pass type I transmembrane glycoprotein 80. Human CD126 is a 468 animo acid singipe pass type I transmembrane glycoprotein 80. Human CD126 artibody, clone B-R6 inhibits IL-6 mediated proliferation of XC-1 cells (transmembrane) glycoprotein 80. Human CD126 artibody, alpha, also known as CD126 IL-R6 or Membrane glycoprotein 80. Human CD126 artibody, and partially blocks binding of IL-6 to its receptor (fourcin et al. 1994), clone B-R6 inhibits IL-6 mediated proliferation of XC-1 cells (transmembrane) glocoprotein 80. Human CD126 Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul Reforences 1. Durandy, A. et al. (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. Jimmunol. 152 (11): 5301-7. 3. Horiuchi, S. et al. (2006) Human T-cell leukenia vitus type-I Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int.J Cancer. 119 (J.H.233). Storage Storage in frost-free freezers is not recommended. Avoid repeated freezing and thaving as this may denature the antibody. Should this product contain a precipitate we recommend microcentifugation before use. Gurarate 12 m	Links	P08887 Related reagents
Fusion Partners Spleen cells from immunised BALB/c (Iffa Credo) mice were fused with the X63/Ag.8653 mouse myeloma cell line. Specificity Mouse anti Human CD126 antibody, clone B-R6 recognizes human Interleukin-6 receptor subunit alpha, also known as CD126, IL-R6 or Membrane glycoprotein 80. Human CD126 is a 468 amino acid single pass type I transmembrane glycoprotein bearing a single Ig-like C2 type domain, two fibronectin type III domains and a single WSXWS motif which may be involved in receptor activation (Dagi <i>et al.</i> 2012). Mouse anti Human CD126 antibody, clone B-R6 inhibits IL-6 mediated proliferation of XG-1 cells (Lu <i>et al.</i> 1995) and partially blocks binding of IL-6 to its receptor (Fourcin <i>et al.</i> 1994), clone B-R6 recognizes both the soluble and membranous human CD126 Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul References 1. Durandy, A. <i>et al.</i> (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. JImmunol. 152 (11): 5361-7. 2. Lu, Z.Y. <i>et al.</i> (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 85 (9): 2251-7. 3. Horiuchi, S. <i>et al.</i> (2009) Ausociation of interleukin-6 signalling with the muscle sexpression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int J Cancer, 119 (4): 823-30. 4. McKay, B.R. <i>et al.</i> (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One, 4: e6027. Storage Storage in frost-free freezers is not reco		
myeloma cell line. Specificity Mouse anti Human CD126 antibody, clone B-R6 recognizes human Interleukin-6 receptor subunit alpha, also known as CD126, IL-6R or Membrane glycoprotein 80. Human CD126 is a 468 amino acid single pass type I transmembrane glycoprotein bearing a single Latike C2 type domain, two fibroneclin type III domains and a single WSXWS motif which may be involved in receptor activation (Dagil <i>et al.</i> 2012). Mouse anti Human CD126 antibody, clone B-R6 inhibits IL-6 mediated proliferation of XG-1 cells (Lu <i>et al.</i> 1995) and partially blocks binding of IL-6 to its receptor (Fourcin <i>et al.</i> 1994), clone B-R6 recognizes both the soluble and membranous human CD126 Removal of Sodium azide is recommended prior to use in functional assays. The use of EOU003 for this purpose. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul References 1. Durandy, A. <i>et al.</i> (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. JImmunol. 152 (11): 5361-7. 2. Lu, Z.Y. <i>et al.</i> (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 36(9): 252-17. 3. Horiuchi, S. <i>et al.</i> (2006) Human T-cell leukemia virus type-I Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int J Cancer. 119 (4): 823-30. 4. McKay, B.R. <i>et al.</i> (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One. 4: e6027. Storage Storage in frost-free freezers	RRID	AB_2127915
alpha, also known as CD126, IL-6R or Membrane glycoprotein 80. Human CD126 is a 468 amino acid single pass type I transmembrane glycoprotein bearing a single Ig-like C2 type domain, two fibronectin type III domains and a single WSXWS motif which may be involved in receptor activation (Dacil <i>et al.</i> 2012). Mouse anti Human CD126 antibody, clone B-R6 inhibits IL-6 mediated proliferation of XG-1 cells (Lu <i>et al.</i> 1995) and partially blocks binding of IL-6 to its receptor (Fourcin <i>et al.</i> 1994), clone B-R6 recognizes both the soluble and membranous human CD126 Removal of Sodium azide is recommended prior to use in functional assays. The use of EQU003 for this purpose. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul References 1. Durandy, A. <i>et al.</i> (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. J Immunol. 152 (11): 5361-7. 2. Lu, Z.Y. <i>et al.</i> (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 85 (9): 2521-7. 3. Horiuchi, S. <i>et al.</i> (2006) Human T-cell leukemia virus type-I Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int J Cancer. 119 (4): 823-30. A. McKay, B.R. <i>et al.</i> (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One. 4: e6027. Storage Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee	Fusion Partners	
(Lu et al. 1995) and partially blocks binding of IL-6 to its receptor (Fourcin et al. 1994), clone B-R6 recognizes both the soluble and membranous human CD126 Removal of Sodium azide is recommended prior to use in functional assays. The use of EQU003 for this purpose. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul References 1. Durandy, A. et al. (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. J Immunol. 152 (11): 5361-7. 2. Lu, Z.Y. et al. (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 85 (9): 2521-7. 3. Horiuchi, S. et al. (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One. 4: e6027. Storage Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/1004.pdf	Specificity	alpha, also known as CD126, IL-6R or Membrane glycoprotein 80. Human CD126 is a 468 amino acid single pass type I transmembrane glycoprotein bearing a single <u>Ig-like C2 type</u> domain, two <u>fibronectin type III</u> domains and a single WSXWS motif which may be involved in receptor
for this purpose. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul References 1. Durandy, A. et al. (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. J Immunol. 152 (11): 5361-7. 2. Lu, Z.Y. et al. (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 85 (9): 2521-7. 3. Horiuchi, S. et al. (2006) Human T-cell leukemia virus type-1 Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int J Cancer. 119 (4): 823-30. 4. McKay, B.R. et al. (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One. 4: e6027. Storage Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf		(Lu et al. 1995) and partially blocks binding of IL-6 to its receptor (Fourcin et al. 1994), clone B-R6
References 1. Durandy, A. et al. (1994) Role of IL-6 in promoting growth of human EBV-induced B-cell tumors in severe combined immunodeficient mice. J Immunol. 152 (11): 5361-7. 2. Lu, Z.Y. et al. (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 85 (9): 2521-7. 3. Horiuchi, S. et al. (2006) Human T-cell leukemia virus type-I Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int J Cancer. 119 (4): 823-30. 4. McKay, B.R. et al. (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One. 4: e6027. Storage Store at +4ºC or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf		
in severe combined immunodeficient mice. J Immunol. 152 (11): 5361-7. 2. Lu, Z.Y. et al. (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. Blood. 85 (9): 2521-7. 3. Horiuchi, S. et al. (2006) Human T-cell leukemia virus type-I Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. Int J Cancer. 119 (4): 823-30. 4. McKay, B.R. et al. (2009) Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. PLoS One. 4: e6027. Storage Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf	Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
This product should be stored undiluted.Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.Guarantee12 months from date of despatchHealth And Safety InformationMaterial Safety Datasheet documentation #10040 available at: 	References	 in severe combined immunodeficient mice. <u>J Immunol. 152 (11): 5361-7.</u> 2. Lu, Z.Y. <i>et al.</i> (1995) Interleukin-10 is a proliferation factor but not a differentiation factor for human myeloma cells. <u>Blood. 85 (9): 2521-7.</u> 3. Horiuchi, S. <i>et al.</i> (2006) Human T-cell leukemia virus type-I Tax induces expression of interleukin-6 receptor (IL-6R): Shedding of soluble IL-6R and activation of STAT3 signaling. <u>Int J Cancer. 119 (4): 823-30.</u> 4. McKay, B.R. <i>et al.</i> (2009) Association of interleukin-6 signalling with the muscle stem cell
Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf	Storage	Store at +4°C or at -20°C if preferred.
may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf		This product should be stored undiluted.
Health And Safety Material Safety Datasheet documentation #10040 available at: Information 10040: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u>		may denature the antibody. Should this product contain a precipitate we recommend
Information 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf	Guarantee	12 months from date of despatch
Regulatory For research purposes only		•
	Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>				
Goat Anti Mouse IgG (STAR77)	HRP			
Rabbit Anti Mouse IgG (STAR12)	RPE			
Rabbit Anti Mouse IgG (STAR8)	DyLight®800			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR76)	RPE			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®680,			
	DI LI LIGAGA FITA LIDD			

DyLight®800, FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

'M373137:200826'

Printed on 26 Aug 2020

© 2020 Bio-Rad Laboratories Inc | Legal | Imprint