

Technical Data Sheet


FITC Mouse Anti-Ki-67 Set

Product Information


Material Number:	556026
Size:	100 Tests
Reactivity:	QC Testing: Human Tested in Development: Mouse Reported Reactivity: Rat, Rhesus
Component:	51-36524X
Description:	FITC Mouse Anti-Ki-67
Size:	100 Tests (1 ea)
Vol. per Test:	20 µl
Clone Name:	B56
Immunogen:	Human Ki-67
Isotype:	Mouse IgG1, κ
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.
Component:	51-35404X
Description:	FITC Mouse IgG1, κ Isotype Control
Size:	100 Tests (1 ea)
Vol. per Test:	20 µl
Clone Name:	MOPC-21
Isotype:	Mouse IgG1, κ
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

Recognizes Ki-67, a nuclear cell proliferation-associated antigen expressed in all active stages of the cell cycle. Ki-67 is revealed as a double band (345 and 395 kDa) in western blot analysis of proliferating cells. B56 was developed using an immunogen composed of the immunodominant epitope of the Ki-67 protein. Antibodies B56 and MIB 1 react with this immunogen. Flow cytometric analysis reveals that the binding of B56-PE can be blocked by MIB 1 purified antibody.



Profile of Ki-67 on MOLT-4 cells analyzed on a FACScan (BDIS, San Jose, CA)



Profile of mouse IgG1 isotype control on MOLT-4 cells analyzed on a FACScan (BDIS, San Jose, CA)

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	866.979.9408	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD

556026 Rev. 8



Application Notes

Application

Intracellular staining (flow cytometry)

Routinely Tested

Recommended Assay Procedure:

Ki-67 staining protocol by flow cytometry:

1. Harvest, count and pellet cells following standard procedures (Note: Ki-67 is expressed by the proliferative cells. You may get no staining with the resting cells, e.g. unstimulated PBMC).
2. While vortexing, add 5 ml drop by drop of cold 70% - 80% ethanol into the cells pellet (1-5 x 10⁷ cells). Then incubate at -20°C for 2 hours minimum. These fixed cells can be used up to 60 days after fixing (Store at -20°C).
3. Add 30-40 ml wash buffer (PBS with 1% FBS, 0.09% NaN₃ pH7.2) to the fixed cells. Centrifuge the cells for 10 minutes at 1000 rpm and aspirate supernatant. Wash one more time with 30-40 ml of wash buffer. Centrifuge at 1000 rpm for 10 minutes and aspirate supernatant.
4. Resuspend the cells to a concentration of 1 X 10⁶/ml (1 X 10⁶/100 µl).
5. Transfer 100 µl cell suspension into each fresh tube.
6. Add 20 µl of properly diluted antibody according to the protocol into the tubes above. Mix gently.
7. Incubate the tubes at room temperature (RT) for 20-30 minutes in the dark.
8. Wash with 2 ml of PBS washing buffer at 1000 rpm for 5 minutes.
9. Aspirate the supernatant.
10. For direct conjugated antibody: go to steps 13 & 14.
11. For purified antibody: add 50 µl of diluted secondary antibody at optimal concentration (Cat. No. 555988), incubate at RT for 30 minutes in the dark.
12. Repeat step 8 & 9.
13. Add 0.5 ml of PBS wash buffer into each tube. For FITC conjugated antibody, add 10 µl of PI Staining Solution (Cat. No. 556463); for PE conjugated antibody, add 20 µl BD Via-Probe™ Cell Viability Solution (Cat. No. 555816) into each tube.
14. Analyze the sample with FACS.

Product Notices

1. This antibody has been optimized and preassayed with its matched isotype control to be used at the recommended volume of 20 ul/test. Titration of the reagents or substituting with other (non-matched) isotype control is NOT recommended.
2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
4. Species testing during development may have been performed with a different format of the same clone. Selected applications have been tested for cross-reactivity.
5. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
6. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

- Benson MJ, Elgueta R, Schpero W, et al. Distinction of the memory B cell response to cognate antigen versus bystander inflammatory signals. *J Exp Med.* 2009; 206(9):2013-2025. (Clone-specific: Flow cytometry)
- Bigley V, Haniffa M, Doulatov S, et al. The human syndrome of dendritic cell, monocyte, B and NK lymphoid deficiency. *J Exp Med.* 2011; 208(2):227-234. (Clone-specific: Flow cytometry)
- Bruno S, Crissman HA, Bauer KD, Darzynkiewicz Z. Changes in cell nuclei during S phase: progressive chromatin condensation and altered expression of the proliferation-associated nuclear proteins Ki-67, cyclin (PCNA), p105, and p34. *Exp Cell Res.* 1991; 196(1):99-106. (Biology: Flow cytometry)
- Bruno S, Darzynkiewicz Z. Cell cycle dependent expression and stability of the nuclear protein detected by Ki-67 antibody in HL-60 cells. *Cell Prolif.* 1992; 25(1):31-40. (Biology: Flow cytometry)
- Kill IR. Localisation of the Ki-67 antigen within the nucleolus: evidence for a fibrillar-deficient region of the dense fibrillar component. *J Cell Sci.* 1996; 109(6):1253-1263. (Biology)
- Kouro T, Medina KL, Oritani K, Kincade PW. Characteristics of early murine B-lymphocyte precursors and their direct sensitivity to negative regulators. *Blood.* 2001; 97(9):2708-2715. (Clone-specific: Flow cytometry)
- Kubbutat MH, Key G, Duchrow M, Schluter C, Flad HD, Gerdes J. Epitope analysis of antibodies recognising the cell proliferation associated nuclear antigen previously defined by the antibody Ki-67 (Ki-67 protein). *J Clin Pathol.* 1994; 47(6):524-528. (Biology)
- Picker LJ, Hagen SI, Lum R, et al. Insufficient production and tissue delivery of CD4+ memory T cells in rapidly progressive simian immunodeficiency virus infection. *J Exp Med.* 2004; 200(10):1299-1314. (Clone-specific: Flow cytometry)
- Pitcher CJ, Hagen SI, Walker JM, et al. Development and homeostasis of T cell memory in rhesus macaque. *J Immunol.* 2002; 168(1):29-43. (Clone-specific: Flow cytometry)
- Scholzen T, Gerdes J. The Ki-67 protein: from the known and the unknown. *J Cell Physiol.* 2000; 182(3):311-322. (Biology)
- Shi SR, Key ME, Kaira KL. Antigen retrieval in formalin-fixed, paraffin-embedded tissues: an enhancement method for immunohistochemical staining based on microwave oven heating of tissue sections. *J Histochem Cytochem.* 1991; 39(6):741-748. (Biology)
- Spargo LDJ, Cleland LG, Cockshell MP, Mayrhofer Graham. Recruitment and proliferation of CD4+ T cells in synovium following adoptive transfer of adjuvant-induced arthritis. *Int Immunol.* 2006; 18(6):897-910. (Clone-specific: Flow cytometry, Immunofluorescence)
- Starborg M, Gell K, Brundell E, Höög C. The murine Ki-67 cell proliferation antigen accumulates in the nucleolar and heterochromatic regions of interphase cells and at the periphery of the mitotic chromosomes in a process essential for cell cycle progression. *J Cell Sci.* 1996; 109(1):143-153. (Biology)
- Valenti LM, Mathieu J, Chancerelle Y, et al. High levels of endogenous nitric oxide produced after burn injury in rats arrest activated T lymphocytes in the first G1 phase of the cell cycle and then induce their apoptosis. *Exp Cell Res.* 2005; 306(1):150-167. (Clone-specific: Flow cytometry)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	866.979.9408	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD

556026 Rev. 8

