Alexa Fluor® 488 anti-human CD127 (IL-7Rα) Antibody

Catalog# / Size	351313 / 25 tests 351314 / 100 tests
Clone	A019D5
Other Names	IL-7 receptor α chain, IL-7Rα
Isotype	Mouse IgG1, κ
Description	CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7Rα. It forms a heterodimer with the common γ chain (γc or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage cells, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be a useful marker for identifying memory and effector T cells. Studies have shown that CD127 expression is down-modulated on Treg cells. It can be used as a marker for differentiation of Treg and conventional T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cell proliferation and development.

Product Details

Reactivity	Human, African Green, Baboon, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Recombinant human CD127
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 488 under optimal conditions.
Concentration	Lot-specific (please contact technical support for concentration and total µg amount, or use our Lookup tool if you have a lot number.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. **Do not freeze.**
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. * Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm. Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.
Excitation Laser	Blue Laser (488 nm)
Application Notes	Additional reported (for the relevant formats) application: proteogenomics[1].
Application References **(PubMed link indicates BioLegend citation)**	1. Peterson VM, et al. 2017. Nat. Biotechnol. 35:936. (PG)

Product Citations **RRID**

1. Azreq M, et al. 2015. J Immunol. 195: 4198 - 4209. PubMed
2. James E, et al. 2016. PLoS Pathog. 12: 1005375. PubMed
3. Abigail E Overacre-Delgoffe et al. 2017. Cell. 169(6):1130-1141 . PubMed
4. van Montfoort N, et al. 2018. Cell. 175:1744. PubMed

AB_10895911 (BioLegend Cat. No. 351313)
AB_10898315 (BioLegend Cat. No. 351314)

Antigen Details

Structure	Type I transmembrane glycoprotein, associates with CD132, 60-90 kD
Distribution	Immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, bone marrow stromal cells
Function	T cell and immature B cell proliferation and development
Ligand/Receptor	IL-7
Cell Type	B cells, T cells, Thymocytes, Tregs
Biology Area	Immunology
Molecular Family	CD Molecules, Cytokine/Chemokine Receptors
Antigen References	1. Sudo T, et al. 1993. P. Natl. Acad. Sci. USA 90:9125. 2. He YW and Malek TR. 1998. Crit. Rev. Immunol. 18:503. 3. Huster KM, et al. 2004. P. Natl. Acad. Sci. USA 101:5610. 4. Pillai M, et al. 2004. Leukemia Lymphoma 45:2403. 5. Morrissey PJ, et al. 1989. J. Exp. Med. 169:707. 6. Liu W, et al. 2006. J. Exp. Med. 203:1701.
Gene ID	3575

Related Protocols

Cell Surface Flow Cytometry Staining Protocol

Other Formats

Purified anti-human CD127 (IL-7Rα), PE anti-human CD127 (IL-7Rα), Pacific Blue™ anti-human CD127 (IL-7Rα), Brilliant Violet 421™ anti-human CD127 (IL-7Rα), FITC anti-human CD127 (IL-7Rα), APC anti-human CD127 (IL-7Rα), Alexa Fluor® 647 anti-human CD127 (IL-7Rα), PE/Cyanine7 anti-human CD127 (IL-7Rα), PerCP/Cyanine5.5 anti-human CD127 (IL-7Rα), Brilliant Violet 570™ anti-human CD127 (IL-7Rα), PE/Cyanine5 anti-human CD127 (IL-7Rα), Brilliant Violet 650™ anti-human CD127 (IL-7Rα), Brilliant Violet 711™ anti-human CD127 (IL-7Rα), Brilliant Violet 785™ anti-human CD127 (IL-7Rα), Brilliant Violet 510™ anti-human CD127 (IL-7Rα), Brilliant Violet 605™ anti-human CD127 (IL-7Rα), PE/Dazzle™ 594 anti-human CD127 (IL-7Rα), Purified anti-human CD127 (IL-7Rα) (Maxpar® Ready), Alexa Fluor® 700 anti-human CD127 (IL-7Rα), Biotin anti-human CD127 (IL-7Rα), APC/Cyanine7 anti-human CD127 (IL-7Rα), APC/Fire™ 750 anti-human CD127 (IL-7Rα), TotalSeq™-A0390 anti-human CD127 (IL-7Rα), TotalSeq™-B0390 anti-human CD127 (IL-7Rα), TotalSeq™-C0390 anti-human CD127 (IL-7Rα), KIRAVIA Blue 520™ anti-human CD127 (IL-7Rα), Spark NIR™ 685 anti-human CD127 (IL-7Rα), PE/Fire™ 640 anti-human CD127 (IL-7Rα)

Product Data



Human peripheral blood lymphocytes were stained with CD3 APC and CD127 (clone A019D5) Alexa Fluor® 488 (top) or mouse IgG1 Alexa Fluor® 488 isotype control (bottom).

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587