

Monoclonal Antibodies Detecting Human Antigens



CD43 (1G10)

Form **Catalog number**
APC-H7 655407

Product availability varies by region. Contact BD Biosciences Customer Support or your local sales representative for information.

RESEARCH APPLICATIONS

Research applications include:

- Research on T cell-macrophage interactions¹
- Research on neutrophil spreading and movement^{2,3}
- Study of *Streptococcus gordonii* DL1 binding to myeloid cells⁴

DESCRIPTION

Specificity

The CD43 antibody reacts specifically with the major 95–135-kilodalton (kDa) sialoglycoprotein found on most human leucocytes.⁵ The CD43 antigen is also known as leukosialin or sialophorin.

Antigen distribution

The CD43 antigen is expressed on T cells, natural killer (NK) cells, pre-B and activated B cells,⁵ granulocytes,⁶ and neutrophils.³ The CD43 antigen is differentially expressed on subpopulations of mononuclear phagocytic cells.⁷ It is not present on most peripheral blood resting B cells, erythrocytes, and non-hematopoietic cells.⁵ Additionally, expression of the CD43 antigen by committed myeloid, erythroid, and lymphoid progenitors has been reported.

The CD43 antigen may be involved in the regulation of B, T, and NK cell function.^{5,8–10} The CD43 antigen plays a role in cell adhesion, which can be positive or negative, depending on the context.^{2,11} The CD43 antigen serves as a counter-receptor for CD169, the prototypical member of the sialic acid binding Ig-like lectins (Siglec) family expressed on macrophages.¹

Clone

The CD43 antibody, clone 1G10,⁵ is derived from the hybridization of X63 mouse myeloma cells with spleen cells isolated from mice immunized with a lymph node suspension from a patient with Hodgkin lymphoma.

Composition

The CD43 antibody is composed of mouse IgG₁ heavy chains and kappa light chains.

Product configuration

The following is supplied in phosphate buffered saline (PBS) containing a stabilizer and a preservative.

Form	Number of tests	Volume per test (μL) ^a	Amount provided (μg)	Total volume (mL)	Concentration (μg/mL)	Stabilizer	Preservative
APC-H7	100	5	25	0.5	50	BSA	ProClin® 300

a. Volume required to stain 10⁶ cells.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Becton, Dickinson and Company
BD Biosciences
2350 Qume Drive
San Jose, CA 95131 USA




CAUTION Some APC-H7 conjugates show changes in their emission spectra with prolonged exposure to paraformaldehyde or light. For overnight storage of stained cells, wash and resuspend in buffer without paraformaldehyde after 1 hour of fixation.

PROCEDURE

Visit our website (bdbiosciences.com) or contact your local BD representative for the lyse/wash protocol for direct immunofluorescence.

REPRESENTATIVE DATA

Flow cytometric analysis was performed on whole blood stained with the indicated conjugated antibody. Laser excitation was at 635 nm. Representative data analyzed with a BD FACS™ brand flow cytometer is shown in the following plot.



HANDLING AND STORAGE

Store vials at 2°C–8°C. Conjugated forms should not be frozen. Protect from exposure to light. Each reagent is stable until the expiration date shown on the bottle label when stored as directed.

WARNING

All biological specimens and materials coming in contact with them are considered biohazards. Handle as if capable of transmitting infection^{12,13} and dispose of with proper precautions in accordance with federal, state, and local regulations. Never pipette by mouth. Wear suitable protective clothing, eyewear, and gloves.

Some reagents are bottled with ProClin 300, and contain 0.003% of a mixture of CMIT/MIT (3:1), CAS number 55965-84-9.



Warning

H317 May cause an allergic skin reaction.

Wear protective gloves/eye protection. Wear protective clothing. Avoid breathing mist/vapours/spray. If skin irritation or rash occurs: Get medical advice/attention. IF ON SKIN: Wash with plenty of water. Dispose of contents/container in accordance with local/regional/national/international regulations.

CHARACTERIZATION

To ensure consistently high-quality reagents, each lot of antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

WARRANTY

Unless otherwise indicated in any applicable BD general conditions of sale for non-US customers, the following warranty applies to the purchase of these products.

THE PRODUCTS SOLD HEREUNDER ARE WARRANTED ONLY TO CONFORM TO THE QUANTITY AND CONTENTS STATED ON THE LABEL OR IN THE PRODUCT LABELING AT THE TIME OF DELIVERY TO THE CUSTOMER. BD DISCLAIMS HEREBY ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR ANY PARTICULAR PURPOSE AND NONINFRINGEMENT. BD'S SOLE LIABILITY IS LIMITED TO EITHER REPLACEMENT OF THE PRODUCTS OR REFUND OF THE PURCHASE PRICE. BD IS NOT LIABLE FOR PROPERTY DAMAGE OR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES, INCLUDING PERSONAL INJURY, OR ECONOMIC LOSS, CAUSED BY THE PRODUCT.

REFERENCES

1. van den Berg TK, Nath D, Ziltener HJ, et al. Cutting edge: CD43 functions as a T cell counterreceptor for the macrophage adhesion receptor sialoadhesin (Siglec-1). *J Immunol.* 2001;166:3637-3640.
2. Matsumoto M, Shigeta A, Miyasaka M, Hirata T. CD43 plays both antiadhesive and proadhesive roles in neutrophil rolling in a context-dependent manner. *J Immunol.* 2008;181:3628-3635.

3. Nathan C, Xie QW, Halbwachs-Mecarelli L, Jin WW. Albumin inhibits neutrophil spreading and hydrogen peroxide release by blocking the shedding of CD43 (sialophorin, leukosialin). *J Cell Biol.* 1993;122:243-256.
4. Urano-Tashiro Y, Yajima A, Takashima E, Takahashi Y, Konishi K. Binding of the Streptococcus gordonii DL1 surface protein Hsa to the host cell membrane glycoproteins CD11b, CD43, and CD50. *Infect Immun.* 2008;76:4686-4691.
5. Horejsi V, Stockinger H. CD43 Workshop Panel report. In: Kishimoto H, Kikutani H, von dem Borne AEGK, et al, eds. *Leucocyte Typing VI: White Cell Differentiation Antigens*. New York: Garland Publishing, Inc.; 1998:494-497.
6. Bazil V, Strominger JL. CD43, the major sialoglycoprotein of human leukocytes, is proteolytically cleaved from the surface of stimulated lymphocytes and granulocytes. *Proc Natl Acad Sci U S A.* 1993;90:3792-3796.
7. Howell DN, Ahuja V, Jones L, Blow O, Sanfilippo FP. Differential expression of CD43 (leukosialin, sialophorin) by mononuclear phagocyte populations. *J Leukoc Biol.* 1994;55:536-544.
8. Bettaieb A, Farace F, Mitjavila MT, et al. Use of a monoclonal antibody (GA3) to demonstrate lineage restricted O-glycosylation on leukosialin during terminal erythroid differentiation. *Blood.* 1988;71:1226-1233.
9. Borche L, Lozano F, Vilella R, Vives J. CD43 monoclonal antibodies recognize the large sialoglycoprotein of human leukocytes. *Eur J Immunol.* 1987;17:1523-1526.
10. Zola H, Swart B, Nicholson I, Voss E. *Leukocyte and Stromal Cell Molecules: The CD Markers*. Hoboken, NJ: John Wiley & Sons, Inc.; 2007.
11. Ardman B, Sikorski MA, Staunton DE. CD43 interferes with T-lymphocyte adhesion. *Proc Natl Acad Sci U S A.* 1992;89:5001-5005.
12. *Protection of Laboratory Workers from Occupationally Acquired Infections; Approved Guideline — Third Edition*. Wayne, PA: Clinical and Laboratory Standards Institute; 2005. CLSI document M29-A3.
13. Centers for Disease Control. Perspectives in disease prevention and health promotion update: universal precautions for prevention of transmission of human immunodeficiency virus, hepatitis B virus, and other bloodborne pathogens in health-care settings. *MMWR.* 1988;37:377-388.

**PATENTS AND
TRADEMARKS**

ProClin is a registered trademark of Rohm and Haas Company.

BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD