

PRODUCT INFORMATION

TasI (Tsp509I)

#ER1351 1000 U

Lot: _____ Expiry Date: __

5'... ↓A A T T ...3'
3'... T T A A↑...5'

Concentration: 10 U/μL
Source: *Thermus aquaticus* Vn4-211
Supplied with: 1 mL of 10X Buffer B
1 mL of 10X Buffer Tango

Store at -20°C



In total 3 vials.

BSA included

www.thermoscientific.com/onebio

RECOMMENDATIONS

1X Buffer B (for 100% TasI digestion)

10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 0.1 mg/mL BSA.

Incubation temperature

65°C*.

Unit Definition

One unit is defined as the amount of TasI required to digest 1 μg of lambda DNA in 1 hour at 65°C in 50 μL of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl, 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol (pH 7.4 at 25°C).

Double Digests

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to the Molecular Biology Tools Product Guide or go to www.thermoscientific.com/doubledigest to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

* Incubate under paraffin oil in a capped vial. Incubation at 37°C results in less than 10% activity.

Storage Buffer

TasI is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/mL BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µL
10X Buffer B	2 µL
DNA (0.5-1 µg/µL)	1 µL
TasI	0.5-2 µL*
- Mix gently and spin down for a few seconds.
- Incubate under paraffin oil in a capped vial at 65°C for 1-16 hours*.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 µL (~0.1-0.5 µg of DNA)
nuclease-free water	18 µL
10X Buffer B	2 µL
TasI	1-2 µL*
- Mix gently and spin down for a few seconds.
- Incubate under paraffin oil in a capped vial at 65°C for 1-16 hours*.

* See Overdigestion Assay on back page.

Thermal Inactivation

TasI is not inactivated by incubation at 80°C for 20 min.

Inactivation Procedure

- To prepare the digested DNA for electrophoresis:
 - stop the digestion reaction by adding 0.5 M EDTA, pH 8.0 (#R1021), to achieve a 20 mM final concentration. Mix thoroughly, add an electrophoresis loading dye and load onto gel.
- To prepare DNA suitable for further enzymatic reactions:
 - extract with phenol/chloroform, precipitate with ethanol or isopropanol, wash the pellet with 75% cold ethanol and air-dry;
 - dissolve DNA in either nuclease-free water, TE buffer, or a buffer suitable for further applications;
 - check the DNA concentration in the solution.

For **ENZYME PROPERTIES** and **CERTIFICATE OF ANALYSIS**
see back page

ENZYME PROPERTIES

Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
100	50-100	20-50	0-20	20-50	0-20

Methylation Effects on Digestion

Dam: never overlaps – no effect.

Dcm: never overlaps – no effect.

CpG: never overlaps – no effect.

EcoKI: never overlaps – no effect.

EcoBI: may overlap – blocked.

Stability during Prolonged Incubation

A minimum of 0.3 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 65°C.

Compatible Ends

EcoRI, MnlI, XapI

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
189	25	8	7	7	10	64

CERTIFICATE OF ANALYSIS

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 80-fold overdigestion with TsaI (5 U/µg lambda DNA x 16 hours).

Ligation and Recleavage (L/R) Assay

The ligation and recleavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of TsaI for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/onebio for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.