



## Contents

Catalog No.	Size
15628019	50 µg
15628050	250 µg

**Kit contents**




## Storage

- Product is shipped at [ambient temperature](#).
- Store at -20°C



## Product description

- The Invitrogen™ 100 bp DNA Ladder is designed for sizing and quantification of double stranded DNA on 1% to 2% agarose gels.
- The 100 bp DNA Ladder consists of 13 individual chromatography-purified DNA fragments ranging in size from 100 bp to 2,000 bp.
- Three reference bands at 2,000 bp, 1,500 bp, and 600 bp are included for easy orientation.
- The ladder is supplied with 10X BlueJuice™ Gel Loading Buffer for sample DNA.



## Important guidelines

- Do not heat the 100 bp DNA Ladder before loading.
- Load the same volume of DNA sample and DNA ladder.
- For quantification, adjust the concentration of the sample to equalize it approximately with the amount of DNA in the nearest band of the ladder.
- For DNA bands visualization with GelRed™ use gel staining after electrophoresis to avoid aberrant DNA migration.



## Guidelines for agarose gel preparation

- Determine the required agarose concentration for your gel based on the size of DNA fragments to be separated.

Fragment size	Recommended agarose gel %	
	1X TAE	1X TBE
800–10,000	0.8	0.7
400–8,000	1.0	0.85
300–7,000	1.2	1.0

- Prepare agarose in a flask with 2-4 times the volume of the agarose solution.
- Exercise caution when handling microwaved agarose. The solution may become superheated and foam over when agitated.
- Refer to the product insert for [UltraPure™ Agarose](#) for detailed instructions on agarose preparation.



## Required materials

**List of materials**

- Visit our [product pages](#) for additional information and protocols.
- Go online to view related [DNA ladders and markers](#).
- For support, visit [thermofisher.com/support](http://thermofisher.com/support).








## Online resources

**Guidelines for staining gels**

**Troubleshooting**

**Limited product warranty and disclaimer details**

## Prepare DNA ladders and samples for electrophoresis

Step	Action												
<p><b>1</b></p> 	<p><b>Cast agarose gel</b></p> <ol style="list-style-type: none"> <li>Prepare agarose solution (w/v) for the gel percentage appropriate for separating your DNA fragments.</li> <li>Microwave agarose solution.</li> <li>Cast agarose gel.</li> </ol>												
<p><b>2</b></p> 	<p><b>Prepare DNA ladder</b></p> <ol style="list-style-type: none"> <li>Thaw, mix and briefly centrifuge each component before use.</li> <li>Add the following components to prepare enough ladder for a single 5 mm well.</li> </ol> <table border="1" data-bbox="976 462 1974 625"> <thead> <tr> <th>Component</th> <th>Volume</th> </tr> </thead> <tbody> <tr> <td>DNA ladder <sup>[1]</sup></td> <td>1 µL (500 ng)</td> </tr> <tr> <td>10X BlueJuice™ Gel Loading Buffer</td> <td>1 µL</td> </tr> <tr> <td>Water, nuclease free</td> <td>8 µL</td> </tr> </tbody> </table> <p>[1] Scale components up or down depending upon width of wells. Modify volume by 0.2 µL (0.1 µg of DNA) for each 1 mm of width.</p> <ol style="list-style-type: none"> <li>Mix gently.</li> <li>Load DNA ladder on gel.</li> </ol>	Component	Volume	DNA ladder <sup>[1]</sup>	1 µL (500 ng)	10X BlueJuice™ Gel Loading Buffer	1 µL	Water, nuclease free	8 µL				
Component	Volume												
DNA ladder <sup>[1]</sup>	1 µL (500 ng)												
10X BlueJuice™ Gel Loading Buffer	1 µL												
Water, nuclease free	8 µL												
<p><b>3</b></p> 	<p><b>Prepare samples</b></p> <ol style="list-style-type: none"> <li>Dilute your sample with 10X BlueJuice™ Gel Loading Buffer (Cat. no. 10816015): mix 1 volume of loading dye with 9 volumes of the DNA sample.</li> <li>Mix gently.</li> <li>Load DNA ladder on gel.</li> </ol>												
<p><b>4</b></p> 	<p><b>Perform electrophoresis</b></p> <ol style="list-style-type: none"> <li>Add appropriate amount of UltraPure TAE or UltraPure TBE buffer to chamber.</li> <li>Set appropriate voltage and perform electrophoresis of samples.</li> </ol> <table border="1" data-bbox="976 1104 1974 1274"> <thead> <tr> <th>DNA size</th> <th>Voltage</th> <th>Buffer</th> </tr> </thead> <tbody> <tr> <td>&lt;1 kb</td> <td>5–10 V/cm</td> <td>TBE</td> </tr> <tr> <td>1–5 kb</td> <td>4–10 V/cm</td> <td>TAE or TBE</td> </tr> <tr> <td>&gt;5 kb</td> <td>1–3 V/cm</td> <td>TAE</td> </tr> </tbody> </table>	DNA size	Voltage	Buffer	<1 kb	5–10 V/cm	TBE	1–5 kb	4–10 V/cm	TAE or TBE	>5 kb	1–3 V/cm	TAE
DNA size	Voltage	Buffer											
<1 kb	5–10 V/cm	TBE											
1–5 kb	4–10 V/cm	TAE or TBE											
>5 kb	1–3 V/cm	TAE											
<p><b>5</b></p> 	<p><b>Stain agarose gel</b></p> <ol style="list-style-type: none"> <li>Incubate gel in staining buffer for 30 minutes.</li> <li>Visualize DNA ladder and samples. <ul style="list-style-type: none"> <li>Use UV transilluminator to detect DNA bands stained with ethidium bromide.</li> <li>Use blue light transilluminator to detect DNA bands stained with SYBR™ stains.</li> </ul> </li> </ol>												